

2/14/00

424 Rec'd PCT/PTO

11 FEB 2000

Express mail label #: EJ918098631US

FORM PTO-1390  
(REV 12-29-99)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

2477US0P

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/485640


INTERNATIONAL APPLICATION NO.  
PCT/JP98/03692

INTERNATIONAL FILING DATE  
August 20, 1998

PRIORITY DATE CLAIMED  
August 21, 1997

TITLE OF INVENTION  
Anti-Inflammatory Agent

APPLICANT(S) FOR DO/EO/US  
H. ODAKA et al.



Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2)) \*
  - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ has been transmitted by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
  - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.  
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

Express mail label #: EJ918098631US

\* This includes specification 31 total pages, including Claims 1-12.

U.S. APPLICATION NO. <b>09/485640</b>		INTERNATIONAL APPLICATION NO. PCT/JP98/03692		ATTORNEY'S DOCKET NUMBER 2477US0P	
---------------------------------------	--	---	--	--------------------------------------	--

17. ☒ The following fees are submitted:

**BASIC NATIONAL FEE ( 37 CFR 1.492 (a) (1) - (5) ) :**

Neither international preliminary examination fee (37 CFR 1.482)  
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO  
and International Search Report not prepared by the EPO or JPO ..... **\$970.00**

International preliminary examination fee (37 CFR 1.482) not paid to  
USPTO but International Search Report prepared by the EPO or JPO ..... **\$840.00**

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but  
international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... **\$690.00**

International preliminary examination fee paid to USPTO (37 CFR 1.482)  
but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... **\$670.00**

International preliminary examination fee paid to USPTO (37 CFR 1.482)  
and all claims satisfied provisions of PCT Article 33(1)-(4) ..... **\$96.00**

**ENTER APPROPRIATE BASIC FEE AMOUNT =**

Surcharge of **\$130.00** for furnishing the oath or declaration later than ☐ 20 ☐ 30  
months from the earliest claimed priority date (37 CFR 1.492(e)).

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	12 - 20 =	0	X \$18.00	\$	0.00
Independent claims	3 - 3 =	0	X \$78.00	\$	0.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$260.00	\$	0.00
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$	840.00

Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement  
must also be filed (Note 37 CFR 1.9, 1.27, 1.28).

**SUBTOTAL =**

Processing fee of **\$130.00** for furnishing the English translation later than ☐ 20 ☐ 30  
months from the earliest claimed priority date (37 CFR 1.492(f)).

**TOTAL NATIONAL FEE =**

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be  
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). **\$40.00** per property +

**TOTAL FEES ENCLOSED =**

	Amount to be refunded:	\$
	charged:	\$

**CALCULATIONS** PTO USE ONLY

a. ☐ A check in the amount of \$\_\_\_\_\_ to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 500799 in the amount of \$ 880.00 to cover the above fees.  
A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any  
overpayment to Deposit Account No. 500799. A duplicate copy of this sheet is enclosed.


  

**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.**

SEND ALL CORRESPONDENCE TO:

Miriam Sohn (Reg. No. 35,368)  
Takeda Pharmaceuticals America, Inc. Patent Office  
1745 Jefferson Davis Highway,  
Suite 408, Arlington VA 22202, USA  
Phone: (703) 416-5160  
Fax: (703) 416-5163  
E-mail: dmsohn@aol.com

SIGNATURE:  
Philippe Y. Riesen

---

NAME  
35,657

---

REGISTRATION NUMBER  
Date: February 7, 2000

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s) : H. ODAKA et al.  
Serial No. : Attn: Box PCT  
Filed on :  
Title : Anti-Inflammatory Agent

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

Preliminary to examination please amend the above-identified application as follows:

IN THE SPECIFICATION:

Page 1, line 30, delete "deivative" and substitute therefor -- derivative--

Page 2, line 13, delete "stimulation was" and substitute therefor --stimulation is--

Page 2, line 30, delete "increased" and substitute therefor -- increases--

Page 22, line 24, delete "wells" and substitute therefor --well--

Page 22, line 28, delete "wells" and substitute therefor --well--

Page 22, line 33, delete "wells" and substitute therefor --well--

Page 22, line 35, delete "wells" and substitute therefor --well--

Page 23, line 2, delete "wells" and substitute therefor --well--

Page 23, line 8, delete "wells" and substitute therefor --well--

Page 23, line 12, delete "wells" and substitute therefor --well--


In the Claims:

Claim 12, page 30, line 23, delete "Use of" and substitute therefor --A method of using--

REMARKS

The above amendments correct typographical and clerical errors and do not constitute new matter. Entry of the above amendments prior to examination is respectfully requested. Early action on the merits is earnestly solicited.

Respectfully submitted,



Philippe Y. Riesen  
Registration No. 35,657  
Attorney for Applicants

c/o Takeda Chemical Industries, Ltd.  
17-85 Jusohonmachi 2-chome  
Yodogawa-ku, Osaka, Japan 532-8686  
Phone #: 81-6-6300-6124  
Fax # : 81-6-6300-6601

Date: February 7, 2000

## DESCRIPTION

ANTI-INFLAMMATORY AGENTTECHNICAL FIELD

- 5       The present invention relates to an anti-inflammatory agent which is useful as an agent for prophylaxis and treatment of a TNF(Tumor Necrosis Factor)- $\alpha$  mediated inflammatory disease.

10   BACKGROUND ART

Regarding a relationship between TNF- $\alpha$  and a thiazolidine derivative, the following references 1) to 4) are known.

- 1) JP-A H7(1995)-285864 describes that a thiazolidine  
15   derivative inhibits production and response reaction of TNF.
- 2) Saishin-Igaku, Vol. 52, No.6, pp.95-102 (1997) describes that a thiazolidine derivative reduces expression of TNF- $\alpha$  and improves insulin-resistance caused by TNF- $\alpha$ .
- 20   3) Endocrinology, Vol. 134, No. 1, pp.264-270 (1994) describes that the overexpression of mRNA for TNF- $\alpha$  and both of its receptors are at least partly normalized by treatment of the diabetic animals with the insulin-sensitizing agent pioglitazone.
- 25   4) Endocrinology, Vol. 136, No. 4, pp.1474-1481 (1995) describes that insulin-sensitizing agents exert their antidiabetic activities by antagonizing the inhibitory effects of TNF- $\alpha$ .

- While, regarding a relationship between an  
30   inflammatory disease and a thiazolidine derivative, the following references 5) and 6) are known.
- 5) WO 96/34943 describes a method for treating a cytokine mediated autoimmune, inflammatory or atherosclerotic disorder with a human 12-lipoxygenase inhibitor. The  
35   human 12-lipoxygenase inhibitor is exemplified by pioglitazone, namely 5-[4-[2-(5-ethyl-2-

pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione.

- 6) The Journal of Biological Chemistry, Vol.271, No.23, pp.13515-13522 (1996) describes that a thiazolidinedione related compound such as 1-(3-allyl-4-oxothiazolidine-2-yliden)-4-methylthiosemicarbazone exhibits antiarthritic activity.

However, none of the above references describes that a thiazolidine derivative is useful as an agent for prophylaxis and treatment of a TNF- $\alpha$  mediated inflammatory disease.

An inflammatory reaction includes various acute and chronic reactions which occur when stimulation was added to the living body. Such reactions include unfavorable reactions which cause destruction of the living tissues as well as favorable reactions to the living body with the purpose of excluding the alien substance. So far, inflammatory diseases are treated with steroid or a nonsteroidal anti-inflammatory agent, an immunosuppressive agent, and the like. However, such agents have problems that they inhibit favorable reactions as well as unfavorable reactions at the time of inflammation.

Therefore, agents which inhibit only unfavorable reactions to the living body are desired.

It is thought that various cytokines are produced to regulate inflammation reactions at the time of inflammation.

TNF- $\alpha$  which is one of such cytokines is thought to play an important role in expansion and delay of inflammation.

For instance, it is thought that production of TNF- $\alpha$  increased to cause destruction of articular tissues in rheumatoid arthritis which belongs to an inflammatory disease.


Based on the above situations, agents which specifically inhibit TNF- $\alpha$  mediated inflammation reactions are expected to be an anti-inflammatory agent with reduced side effects, therefore development of such

agents are desired.

# DISCLOSURE OF INVENTION

The present invention relates to

- 5 (1) An anti-inflammatory agent which affects by way of a TNF- $\alpha$  inhibitory action and comprises a compound of the formula:



- wherein R represents a hydrocarbon group that may be substituted or a heterocyclic group that may be substituted; Y represents a group of the formula -CO-, -CH(OH)-, or -NR<sup>3</sup>- where R<sup>3</sup> represents an alkyl group that may be substituted; m is 0 or 1; n is 0, 1 or 2; X represents CH or N; A represents a chemical bond or a bivalent aliphatic hydrocarbon group having 1 to 7 carbon atoms; Q represents oxygen or sulfur; R<sup>1</sup> represents hydrogen or an alkyl group; ring E may have further 1 to 4 substituents, which may form a ring in combination with R<sup>1</sup>; L and M respectively represent hydrogen or may be combined with each other to form a chemical bond; or a salt thereof (hereinafter referred to simply as Compound (I));
- 15
- 20
- 25

- (2) An anti-inflammatory agent according to the above (1), wherein the heterocyclic group represented by R is a 5- to 7-membered monocyclic and heterocyclic group containing 1 to 4 hetero-atoms selected from oxygen, sulfur and nitrogen in addition to carbon as ring members or a condensed heterocyclic group;
- 30


- (3) An anti-inflammatory agent according to the above (1), wherein R represents a heterocyclic group that may be substituted;
- 35

- (4) An anti-inflammatory agent according to the above (3),

wherein the heterocyclic group is pyridyl, oxazolyl or thiazolyl;

(5) An anti-inflammatory agent according to the above (1), wherein the partial structural formula:

5



(6) An anti-inflammatory agent according to the above (1), wherein X represents CH;

(7) An anti-inflammatory agent according to the above (1), wherein R<sup>1</sup> represents hydrogen;

(8) An anti-inflammatory agent according to the above (1), wherein L and M respectively represent hydrogen;

(9) An anti-inflammatory agent according to the above (1), wherein the compound is 5-[4-[2-(5-ethyl-2-pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione;

(10) An anti-inflammatory agent according to the above (1), wherein the compound is (R)-(+)-5-[3-[4-[2-(2-furyl)-5-methyl-4-oxazolylmethoxy]-3-methoxyphenyl]propyl]-2,4-oxazolidinedione;

(11) Method for treating or preventing a TNF- $\alpha$  mediated inflammatory disease in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound as defined in the above (1) or a pharmacologically acceptable salt thereof; and

(12) Use of a compound as defined in the above (1) or a pharmacologically acceptable salt thereof for the manufacture of an agent for prophylaxis or treatment of a TNF- $\alpha$  mediated inflammatory disease.

Referring to the hydrocarbon group that may be substituted for R, the hydrocarbon group includes aliphatic, alicyclic, alicyclic-aliphatic, aromatic-aliphatic, and aromatic hydrocarbon groups. The number of carbon atoms constituting such hydrocarbon groups is preferably 1 to 14.



The aliphatic hydrocarbon group is preferably a  $C_{1-8}$  aliphatic hydrocarbon group. The aliphatic hydrocarbon group includes saturated  $C_{1-8}$  aliphatic hydrocarbon groups (e.g. alkyl groups) such as methyl, ethyl, propyl, 5 isopropyl, butyl, isobutyl, sec-butyl, t-butyl, pentyl, isopentyl, neopentyl, t-pentyl, hexyl, isohexyl, heptyl, and octyl; and unsaturated  $C_{2-8}$  aliphatic hydrocarbon groups (e.g. alkenyl, alkadienyl, alkynyl, and alkadiynyl groups) such as ethenyl, 1-propenyl, 2-propenyl, 1-butenyl, 2- 10 butenyl, 3-butenyl, 2-methyl-1-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 3-methyl-2-butenyl, 1-hexenyl, 3-hexenyl, 2,4-hexadienyl, 5-hexenyl, 1-heptenyl, 1-octenyl, ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 3-butynyl, 1-pentynyl, 2-pentynyl, 15 3-pentynyl, 4-pentynyl, 1-hexynyl, 3-hexynyl, 2,4-hexadiynyl, 5-hexynyl, 1-heptynyl, and 1-octynyl.

The alicyclic hydrocarbon group is preferably a  $C_{3-7}$  alicyclic hydrocarbon group. The alicyclic hydrocarbon group includes saturated  $C_{3-7}$  alicyclic hydrocarbon groups 20 (e.g. cycloalkyl groups) such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, etc. and unsaturated  $C_{5-7}$  alicyclic hydrocarbon groups (e.g. cycloalkenyl groups and cycloalkadienyl groups) such as 1-cyclopentenyl, 2-cyclopentenyl, 3-cyclopentenyl, 1-cyclohexenyl, 2- 25 cyclohexenyl, 3-cyclohexenyl, 1-cycloheptenyl, 2-cycloheptenyl, 3-cycloheptenyl, and 2,4-cycloheptadienyl.

The alicyclic-aliphatic hydrocarbon group is a group consisting of the above-described alicyclic hydrocarbon group and aliphatic hydrocarbon group (e.g. cycloalkyl- 30 alkyl and cycloalkenyl-alkyl groups) and is preferably a  $C_{4-9}$  alicyclic-aliphatic hydrocarbon group. Specifically, the alicyclic-aliphatic hydrocarbon group includes cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclopentylmethyl, 2-cyclopentenylmethyl, 3- 35 cyclopentenylmethyl, cyclohexylmethyl, 2-

cyclohexenylmethyl, 3-cyclohexenylmethyl, cyclohexylethyl, cyclohexylpropyl, cycloheptylmethyl, cycloheptylethyl, etc.

The aromatic-aliphatic hydrocarbon group is preferably a C<sub>7-13</sub> aromatic-aliphatic hydrocarbon group (e.g. aralkyl and aryl-alkenyl groups). The aromatic-aliphatic hydrocarbon group includes C<sub>7-</sub>, phenylalkyl such as benzyl, phenethyl, 1-phenylethyl, 3-phenylpropyl, 2-phenylpropyl and 1-phenylpropyl; C<sub>11-13</sub> naphthylalkyl such as  $\alpha$ -naphthylmethyl,  $\alpha$ -naphthylethyl,  $\beta$ -naphthylmethyl, and  $\beta$ -naphthylethyl; C<sub>8-10</sub> phenylalkenyl such as styryl and 4-phenyl-1,3-butadienyl; and C<sub>12-13</sub> naphthylalkenyl such as 2-(2-naphthyl)vinyl.

The aromatic hydrocarbon group is preferably a C<sub>6-14</sub> aromatic hydrocarbon group (e.g. aryl groups). The aromatic hydrocarbon group includes phenyl and naphthyl ( $\alpha$ -naphthyl,  $\beta$ -naphthyl).

Referring to the formula (I), the heterocyclic group in a heterocyclic group that may be substituted for R is a 5- to 7-membered monocyclic and heterocyclic group containing 1 to 4 hetero-atoms selected from oxygen, sulfur, and nitrogen in addition to carbon as ring members or a condensed heterocyclic group. The condensed heterocyclic group may for example be one consisting of such a 5- to 7-membered monocyclic and heterocyclic group and a 6-membered ring containing 1 or 2 nitrogen atoms, a benzene ring, or a 5-membered ring containing one sulfur atom.

Specifically the heterocyclic group includes 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrazinyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, isothiazolyl, isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 1,2,4-oxadiazol-5-yl, 1,2,4-triazol-3-yl, 1,2,3-triazol-4-yl, tetrazol-5-yl, benzimidazol-2-yl,

indol-3-yl, 1H-indazol-3-yl, 1H-pyrrolo[2,3-b]pyrazin-2-yl, 1H-pyrrolo[2,3-b]pyridin-6-yl, 1H-imidazo[4,5-b]pyridin-2-yl, 1H-imidazo[4,5-c]pyridin-2-yl, 1H-imidazo[4,5-b]pyrazin-2-yl, benzopyranyl and 3,4-

- 5 dihydrobenzopyran-2-yl. The preferred heterocyclic group is pyridyl, oxazolyl, or thiazolyl.

Referring to the formula (I), the hydrocarbon group and heterocyclic group for R may respectively have 1 to 5, preferably 1 to 3 substituents at substitutable positions.

- 10 Such substituents include for example aliphatic hydrocarbon groups, alicyclic hydrocarbon groups, aryl groups, aromatic heterocyclic groups, non-aromatic heterocyclic groups, halogen, nitro, amino group that may be substituted, acyl groups that may be substituted,  
15 hydroxy group that may be substituted, thiol that may be substituted, and carboxyl group that may be esterified.

- The aliphatic hydrocarbon group includes straight-chain or branched aliphatic hydrocarbon groups having 1 to 15 carbon atoms, such as alkyl groups, alkenyl groups, and  
20 alkynyl groups.

- The preferred alkyl group is a  $C_{1-10}$  alkyl group, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, t-butyl, pentyl, isopentyl, neopentyl, t-pentyl, 1-ethylpropyl, hexyl, isohexyl, 1,1-dimethylbutyl, 2,2-  
25 dimethylbutyl, 3,3-dimethylbutyl, 2-ethylbutyl, hexyl, pentyl, octyl, nonyl, and decyl.

- The preferred alkenyl group is a  $C_{2-10}$  alkenyl group, such as vinyl, allyl, isopropenyl, 1-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 2-  
30 ethyl-1-butenyl, 3-methyl-2-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 4-methyl-3-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, and 5-hexenyl.

- The preferred alkynyl group is a  $C_{2-10}$  alkynyl group, such as ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butynyl, 3-butynyl, 1-pentyne, 2-pentyne, 3-pentyne, 4-pentyne, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl,  
35

and 5-hexynyl.

The alicyclic hydrocarbon group includes saturated and unsaturated alicyclic hydrocarbon groups having 3 to 12 carbon atoms, such as cycloalkyl groups, cycloalkenyl groups, and cycloalkadienyl groups.

The preferred cycloalkyl group is a  $C_{3-10}$  cycloalkyl group, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl, bicyclo[3.2.1]octyl, bicyclo[3.2.2]nonyl, bicyclo[3.3.1]nonyl, bicyclo[4.2.1]nonyl, and bicyclo[4.3.1]decyl.

The preferred cycloalkenyl group is a  $C_{3-10}$  cycloalkenyl group, such as 2-cyclopenten-1-yl, 3-cyclopenten-1-yl, 2-cyclohexen-1-yl, and 3-cyclohexen-1-yl.

The preferred cycloalkadienyl group is a  $C_{4-10}$  cycloalkadienyl group, such as 2,4-cyclopentadien-1-yl, 2,4-cyclohexadien-1-yl, 2,5-cyclohexadien-1-yl.

The term "aryl group" means a monocyclic or condensed polycyclic aromatic hydrocarbon group. As preferred examples,  $C_{6-14}$  aryl groups such as phenyl, naphthyl, anthryl, phenanthryl, acenaphthylenyl can be mentioned. Particularly preferred are phenyl, 1-naphthyl, and 2-naphthyl.

The preferred aromatic heterocyclic group includes 5- to 7-membered monocyclic aromatic heterocyclic groups containing 1 to 4 hetero-atoms selected from oxygen, sulfur, and nitrogen in addition to carbon as ring members, such as furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, furazanyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, tetrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl; and bicyclic or tricyclic condensed aromatic

heterocyclic groups containing 1 to 5 hetero-atoms selected from oxygen, sulfur, and nitrogen in addition to carbon as ring members, such as benzofuranyl, isobenzofuranyl, benzo[b]thienyl, indolyl, isoindolyl, 1H-indazolyl, 5 benzimidazolyl, benzoxazolyl, 1,2-benzisoxazolyl, benzothiazolyl, 1,2-benzisothiazolyl, 1H-benzotriazolyl, quinolyl, isoquinolyl, cinnolinyl, quinazolinyl, quinoxalinyl, phthalazinyl, naphthyridinyl, purinyl, pteridinyl, carbazolyl,  $\alpha$ -carbolinyl,  $\beta$ -carbolinyl,  $\gamma$  10 -carbolinyl, acridinyl, phenoxazinyl, phenothiazinyl, phenazinyl, phenoxathiinyl, thianthrenyl, phenanthridinyl, phenanthrolinyl, indolizinyl, pyrrolo[1,2-b]pyridazinyl, pyrazolo[1,5-a]pyridyl, imidazo[1,2-a]pyridyl, imidazo[1,5-a]pyridyl, 15 imidazo[1,2-b]pyridazinyl, imidazo[1,2-a]pyrimidinyl, 1,2,4-triazolo[4,3-a]pyridyl, and 1,2,4-triazolo[4,3-b]pyridazinyl.

The preferred non-aromatic heterocyclic group includes oxiranyl, azetidiny, oxetanyl, thietanyl, 20 pyrrolidinyl, tetrahydrofuryl, thiolanyl, piperidyl, tetrahydropyranyl, morpholinyl, thiomorpholinyl, piperazinyl, pyrrolidino, piperidino, and morpholino.

The halogen includes fluorine, chlorine, bromine, and iodine, and is preferably fluorine or chlorine.

25 The amino group that may be substituted includes amino ( $-\text{NH}_2$ ) that may be mono- or di-substituted by, for example,  $\text{C}_{1-10}$  alkyl groups,  $\text{C}_{3-10}$  cycloalkyl groups,  $\text{C}_{2-10}$  alkenyl groups,  $\text{C}_{3-10}$  cycloalkenyl groups,  $\text{C}_{1-13}$  acyl groups (e.g.  $\text{C}_{2-10}$  alkanoyl groups,  $\text{C}_{7-13}$  arylcarbonyl groups), or  $\text{C}_{6-12}$  aryl 30 groups. As examples of the substituted amino group, there can be mentioned methylamino, dimethylamino, ethylamino, diethylamino, dibutylamino, diallylamino, cyclohexylamino, acetylamino, propionylamino, benzoylamino, phenylamino, and N-methyl-N-phenylamino.

35 The acyl group in the acyl groups that may be substituted includes  $\text{C}_{1-13}$  acyl groups. For example, formyl

and groups formed between carbonyl and  $C_{1-10}$  alkyl groups,  $C_{3-10}$  cycloalkyl groups,  $C_{2-10}$  alkenyl groups,  $C_{3-10}$  cycloalkenyl groups,  $C_{6-12}$  aryl groups, or aromatic heterocyclic groups (e.g. thienyl, furyl, pyridyl). The preferred acyl group includes acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, heptanoyl, octanoyl, cyclobutanecarbonyl, cyclopentanecarbonyl, cyclohexanecarbonyl, cycloheptanecarbonyl, crotonyl, 2-cyclohexenecarbonyl, benzoyl, and nicotinoyl. The substituent in the substituted acyl groups includes  $C_{1-3}$  alkyl,  $C_{1-3}$  alkoxy groups, halogen (e.g. chlorine, fluorine, bromine, etc.), nitro, hydroxy, and amino.

Referring to the hydroxy group that may be substituted, the substituted hydroxy includes alkoxy, alkenyloxy, aralkyloxy, acyloxy, and aryloxy groups.

The preferred alkoxy group includes  $C_{1-10}$  alkoxy groups, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, t-butoxy, pentyloxy, isopentyloxy, neopentyloxy, hexyloxy, heptyloxy, nonyloxy, cyclobutoxy, cyclopentyloxy, and cyclohexyloxy.

The preferred alkenyloxy group includes  $C_{2-10}$  alkenyloxy groups, such as allyloxy, crotyloxy, 2-pentenyl, 3-hexenyl, 2-cyclopentenylmethoxy, and 2-cyclohexenylmethoxy.

The preferred aralkyloxy group includes  $C_{7-10}$  aralkyloxy groups, such as phenyl- $C_{1-4}$  alkyloxy (e.g. benzyloxy, phenethyloxy, etc.).

The preferred acyloxy group includes  $C_{2-13}$  acyloxy groups, more preferably  $C_{2-4}$  alkanoyloxy (e.g. acetyloxy, propionyloxy, butyryloxy, isobutyryloxy, etc.).

The preferred aryloxy group includes  $C_{6-14}$  aryloxy groups, such as phenoxy, and naphthyloxy. This aryloxy group may have 1 or 2 substituents such as halogen (e.g. chlorine, fluorine, bromine, etc.). The substituted aryloxy group includes 4-chlorophenoxy.

Referring to the thiol group that may be substituted, the substituted thiol group includes alkylthio, cycloalkylthio, aralkylthio, and acylthio groups.

The preferred alkylthio group includes  $C_{1-10}$  alkylthio groups, such as methylthio, ethylthio, propylthio, isopropylthio, butylthio, isobutylthio, sec-butylthio, t-butylthio, pentylthio, isopentylthio, neopentylthio, hexylthio, heptylthio, and nonylthio. The preferred cycloalkylthio group includes  $C_{3-10}$  cycloalkylthio groups such as cyclobutylthio, cyclopentylthio, and cyclohexylthio.

The preferred aralkylthio group includes  $C_{7-10}$  aralkylthio groups, such as phenyl- $C_{1-4}$  alkylthio (e.g. benzylthio, phenethylthio, etc.).

The acylthio group is preferably a  $C_{2-13}$  acylthio group, more preferably a  $C_{2-4}$  alkanoylthio group (e.g. acetylthio, propionylthio, butyrylthio, isobutyrylthio, etc.).

The carboxyl group that may be esterified includes alkoxycarbonyl, aralkyloxycarbonyl, and aryloxycarbonyl groups.

The preferred alkoxycarbonyl group includes  $C_{2-5}$  alkoxycarbonyl groups, such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, and butoxycarbonyl.

The preferred aralkyloxycarbonyl group includes  $C_{8-10}$  aralkyloxycarbonyl groups, such as benzyloxycarbonyl.

The preferred aryloxycarbonyl group includes  $C_{7-15}$  aryloxycarbonyl groups, such as phenoxycarbonyl, and p-tolyloxycarbonyl.

The preferred substituent on the hydrocarbon or heterocyclic group for R includes  $C_{1-10}$  alkyl groups, aromatic heterocyclic groups, and  $C_{6-14}$  aryl groups. Particularly preferred is  $C_{1-3}$  alkyl, furyl, thienyl, phenyl, or naphthyl.

Referring to the formula (I), when the substituent on the hydrocarbon or heterocyclic group for R is an alicyclic hydrocarbon group, an aryl group, an aromatic heterocyclic

group, or a non-aromatic heterocyclic group, this substituent may be further substituted by one or more, preferably 1 to 3 suitable substituents. As such substituents, there can be mentioned  $C_{1-6}$  alkyl groups,  $C_{2-6}$  alkenyl groups,  $C_{2-6}$  alkynyl groups,  $C_{3-7}$  cycloalkyl groups,  $C_{6-14}$  aryl groups (e.g. phenyl, naphthyl, etc.), aromatic heterocyclic groups (e.g. thienyl, furyl, pyridyl, oxazolyl, thiazolyl, etc.), non-aromatic heterocyclic groups (e.g. tetrahydrofuryl, morpholino, thiomorpholino, piperidino, pyrrolidino, piperazino, etc.),  $C_{7-9}$  aralkyl groups, amino, N-mono( $C_{1-4}$ )alkylamino groups, N,N-di( $C_{1-4}$ )alkylamino groups,  $C_{2-8}$  acylamino groups (e.g. acetylamino, propionylamino, benzoylamino, etc.), amidino,  $C_{2-8}$  acyl groups (e.g.  $C_{2-8}$  alkanoyl groups, etc.), carbamoyl, N-mono( $C_{1-4}$ )alkylcarbamoyl groups, N,N-di( $C_{1-4}$ )alkylcarbamoyl groups, sulfamoyl, N-mono( $C_{1-4}$ )alkylsulfamoyl groups, N,N-di( $C_{1-4}$ )alkylsulfamoyl groups, carboxyl,  $C_{2-8}$  alkoxy carbonyl groups, hydroxy,  $C_{1-4}$  alkoxy groups,  $C_{2-5}$  alkenyloxy groups,  $C_{3-7}$  cycloalkyloxy groups,  $C_{7-9}$  aralkyloxy groups,  $C_{6-14}$  aryloxy groups (e.g. phenyloxy, naphthyloxy, etc.), mercapto,  $C_{1-4}$  alkylthio groups,  $C_{7-9}$  aralkylthio groups,  $C_{6-14}$  arylthio groups (e.g. phenylthio, naphthylthio, etc.), sulfo, cyano, azido, nitro, nitroso, and halogen (e.g. fluorine, chlorine, bromine, iodine).

In the formula (I), R is preferably a heterocyclic group that may be substituted. More preferably, R is pyridyl, oxazolyl, or thiazolyl group, which may have 1 to 3 substituents selected from  $C_{1-3}$  alkyl, furyl, thienyl, phenyl, and naphthyl.

Referring to the formula (I), Y represents  $-CO-$ ,  $-CH(OH)-$  or  $-NR^3-$ . Y is preferably  $-CH(OH)-$  or  $-NR^3-$  and more preferably  $-CH(OH)-$ . Referring to an alkyl group that may be substituted for  $R^3$ , the alkyl group includes  $C_{1-4}$  alkyl groups, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, and t-butyl. The substituent includes halogen (e.g. fluorine, chlorine, bromine,



iodine),  $C_{1-4}$  alkoxy groups (e.g. methoxy, ethoxy, propoxy, butoxy, isobutoxy, sec-butoxy, t-butoxy), hydroxy, nitro, and  $C_{1-4}$  acyl groups (e.g. formyl, acetyl, propionyl, etc.).

The symbol m represents 0 or 1 and is preferably 0.



5 The symbol n represents 0, 1 or 2 and is preferably 0 or 1.

The symbol X represents CH or N and is preferably CH.

Referring to the formula (I), the symbol A represents a chemical bond or a bivalent aliphatic hydrocarbon group having 1 to 7 carbon atoms. This aliphatic hydrocarbon group may be straight-chain or branched and may further be saturated or unsaturated. Thus, for example,  $-CH_2-$ ,  $-CH(CH_3)-$ ,  $-(CH_2)_2-$ ,  $-CH(C_2H_5)-$ ,  $-(CH_2)_3-$ ,  $-(CH_2)_4-$ ,  $-(CH_2)_5-$ ,  $-(CH_2)_6-$ ,  $-(CH_2)_7-$ , etc. can be mentioned for the saturated bivalent aliphatic hydrocarbon group, while  $-CH=CH-$ ,  $-C(CH_3)=CH-$ ,  $-CH=CH-CH_2-$ ,  $-C(C_2H_5)=CH-$ ,  $-CH_2-CH=CH-CH_2-$ ,  $-CH_2-CH_2-CH=CH-CH_2-$ ,  $-CH=CH-CH=CH-CH_2-$ ,  $-CH=CH-CH=CH-CH=CH-CH_2-$ , etc. can be mentioned for the unsaturated bivalent aliphatic hydrocarbon group. The symbol A preferably represents a chemical bond or a bivalent aliphatic hydrocarbon group having 1 to 4 carbon atoms, which is preferably a saturated group. More preferably, A represents a chemical bond,  $-CH_2-$  or  $-(CH_2)_2-$ . Still more preferably, A represents a chemical bond or  $-(CH_2)_2-$ .

25 The alkyl group for  $R^1$  includes  $C_{1-4}$  alkyl groups such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, and t-butyl. Preferably,  $R^1$  represents hydrogen.

Referring to the formula (I), the partial structural formula:




35 wherein each symbols has the same meanings as defined above.

Furthermore, ring E may optionally have 1 to 4

substituents at substitutable positions. Such substituents include an alkyl group, a hydroxy group that may be substituted, halogen, an acyl group that may be substituted, nitro, and an amino group that may be substituted. These substituents may be the same as the substituents mentioned for the hydrocarbon or heterocyclic group for R.

Ring E, the partial structural formula:




wherein  $R^2$  represents hydrogen, an alkyl group, a hydroxy group that may be substituted, halogen, an acyl group that may be substituted, nitro, or an amino group that may be substituted.

The alkyl group, hydroxy group that may be substituted, halogen, acyl group that may be substituted, and amino group that may be substituted, for  $R^2$ , may each be the same as the substituents mentioned for the hydrocarbon or heterocyclic group for R. Preferably,  $R^2$  is hydrogen, hydroxy group that may be substituted, or halogen. More preferably,  $R^2$  is hydrogen or hydroxy group that may be substituted. Particularly preferred is hydrogen or a  $C_{1-4}$  alkoxy group.


L and M respectively represent hydrogen or may be combined with each other to form a chemical bond, and preferably they are hydrogen.

Referring to the formula (I), the compound in which L and M are combined with each other to form a chemical bond:



wherein each symbols has the same meanings as defined above, may exist as (E)- and (Z)- isomers, owing to the double bond at 5-position of the azolidinedione ring.


The compound in which L and M respectively represent  
5 hydrogen:



10

wherein each symbols has the meanings as defined above, may exist as optical isomers, i.e. (R)- and (S)-forms, with respect to the asymmetric carbon at 5-position of the  
15 azolidinedione ring. This compound includes those optically active compounds, i.e. (R)- and (S)-forms, as well as the racemic form.

The preferred compound of the formula (I) is the compound in which R represents pyridyl, oxazolyl, or  
20 thiazolyl group, optionally having 1 to 3 substituents selected from the group consisting of C<sub>1-3</sub> alkyl, furyl, thienyl, phenyl, and naphthyl; Y represents -CH(OH)- or -NR<sup>3</sup>- wherein R<sup>3</sup> is methyl; n is 0 or 1; X represents CH; A represents a chemical bond or -(CH<sub>2</sub>)<sub>2</sub>-; R<sup>1</sup> represents  
25 hydrogen; ring E, namely the partial structural formula:



30 wherein R<sup>2</sup> is hydrogen or a C<sub>1-4</sub> alkoxy group; and L and M respectively represent hydrogen.

As preferred species of the compound of the formula (I), the following compounds are mentioned.

- 35 1) 5-[4-[2-(5-ethyl-2-pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione;  
2) 5-[4-[2-hydroxy-2-(5-methyl-2-phenyl-4-

oxazolyl)ethoxy]benzyl]-2,4-thiazolidinedione;

3) (R)-(+)-5-[3-[4-[2-(2-furyl)-5-methyl-4-oxazolylmethoxy]-3-methoxyphenyl]propyl]-2,4-oxazolidinedione;

5 4) (S)-(-)-5-[3-[4-[2-(2-furyl)-5-methyl-4-oxazolylmethoxy]-3-methoxyphenyl]propyl]-2,4-oxazolidinedione;

5) 5-[3-[3-fluoro-4-(5-methyl-2-phenyl-4-oxazolylmethoxy)phenyl]propyl]-2,4-oxazolidinedione;

10 6) 5-[5-[3-methoxy-4-(5-methyl-2-phenyl-4-oxazolylmethoxy)phenyl]pentyl]-2,4-oxazolidinedione;

7) 5-[3-[3,5-dimethoxy-4-[2-[(E)-styryl]-4-oxazolylmethoxy]phenyl]propyl]-2,4-oxazolidinedione;

8) 5-[[4-[(3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl]methyl]-2,4-thiazolidinedione;

9) 5-[[4-[2-(methyl-2-pyridylamino)ethoxy]phenyl]methyl]-2,4-thiazolidinedione.

20 Hereafter, these compounds are sometimes simply referred to as compound No.1, compound No.2, and the like.

Among the above compounds, compound Nos. 1, 3, 8 and 9 are preferred, and compound Nos.1 and 3 are particularly preferred.

25 The salt of compound (I) of the present invention is preferably a pharmacologically acceptable salt, which includes salts with inorganic bases, salts with organic bases, salts with inorganic acids, salts with organic acids, and salts with basic or acidic amino acids.

30 The preferred salt with an inorganic base includes alkali metal salts such as sodium salt, potassium salt, etc.; alkaline earth metal salts such as calcium salt, magnesium salt, etc.; aluminum salt, and ammonium salts.

35 The preferred salt with an organic base includes salts with trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine,

dicyclohexylamine, N,N'-dibenzylethylenediamine, etc.

The preferred salt with an inorganic acid includes salts with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, etc.

5       The preferred salt with an organic acid includes salts with formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, etc.

10       The preferred salt with a basic amino acid includes salts with arginine, lysine, ornithine, etc. The preferred salt with an acidic amino acid includes salts with aspartic acid, glutamic acid, etc.

15       The most preferred of all the above-mentioned salts is hydrochloride, sodium salt or potassium salt.

20       Compound (I) or a salt thereof of the present invention can be produced in accordance with methods described in JP-A S55(1980)-22636 (EP-A-8203), JP-A S60(1985)-208980 (EP-A-155845), JP-A S61(1986)-286376 (EP-A-208420), JP-A S61(1986)-085372 (EP-A-177353), JP-A S61(1986)-267580 (EP-A-193256), JP-A H5(1993)-86057 (WO-A-9218501), JP-A H7(1995)-82269 (EP-A-605228), JP-A H7(1995)-101945 (EP-A-612743), EP-A-643050, EP-A-710659 (JP-A H9(1997)-194467), etc, or methods analogous thereto.

25

30       Compound (I) or a salt thereof of the present invention (hereinafter simply referred to as compound of the present invention) is useful as an anti-inflammatory agent which affects by way of a TNF- $\alpha$  inhibitory action. In addition, the toxic potential of the compound of the present invention is low. The TNF- $\alpha$  inhibitory action means reduction in the production amount of TNF- $\alpha$  in the living tissues (e.g., skeletal muscles, monocytes, macrophages, neutrophils, fibroblasts, epithelial cells, astrocytes, etc.) and  
35       reduction in the activity of TNF- $\alpha$ .

The anti-inflammatory agent of the present invention can be used as an agent for prophylaxis and treatment of TNF- $\alpha$  mediated inflammatory diseases in mammals (e.g., man, mouse, rat, rabbit, dog, cat, bovine, equine, swine, monkey, etc.). The TNF- $\alpha$  mediated inflammatory diseases mean inflammatory diseases which occur in the presence of TNF- $\alpha$  and can be treated by way of a TNF- $\alpha$  inhibitory action.

Examples of such inflammatory diseases include diabetic complications (e.g., retinopathy, nephropathy, neuropathy, disorders in the great arteries, etc.), rheumatoid arthritis, osteoarthritis of the spine, osteoarthritis, low back pain, gout, postoperative or traumatic inflammation, remission of swelling, neuralgia, laryngopharyngitis, cystitis, hepatitis, pneumonia, etc.

As the anti-inflammatory agent of the present invention, the compound of the present invention as such can be used. Usually, the anti-inflammatory agent is used in the form of a pharmaceutical composition obtained by formulating the compound of the invention with per se known pharmaceutically acceptable carriers.

As the pharmaceutically acceptable carrier, a variety of organic and inorganic carriers in common use as raw materials for pharmaceutical preparations are employed. The carrier is formulated in the form of the excipient, lubricant, binder, and disintegrator for a solid dosage form; and the solvent, solubilizer, suspending agent, isotonizing agent, buffering agent and local analgesic for a liquid dosage form. When necessary, pharmaceutical additives such as the preservative, antioxidant, coloring agent, sweetener, etc. can also be used.

The preferred excipient includes lactose, sucrose, D-mannitol, starch, crystalline cellulose, light silicic anhydride, etc.

The preferred lubricant includes magnesium stearate, calcium stearate, talc, colloidal silica, etc.

The preferred binder includes crystalline cellulose, sucrose, D-mannitol, trehalose, dextrin, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, etc.

- 5       The preferred disintegrator includes starch, carboxymethylcellulose, carboxymethylcellulose calcium, croscarmellose sodium, carboxymethylstarch sodium, etc.

10       The preferred solvent includes water for injection, alcohol, propylene glycol, macrogol, sesame oil, corn oil, tricaprylin, etc.

      The preferred solubilizer includes polyethylene glycol, propylene glycol, D-mannitol, trehalose, benzyl benzoate, ethanol, trisaminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate, etc.

- 15       The preferred suspending agent includes surfactants such as stearyltriethanolamine, sodium lauryl sulfate, laurylaminopropionic acid, lecithin, benzalkonium chloride, benzethonium chloride, glyceryl monostearate, etc. and hydrophilic polymers such as polyvinyl alcohol,  
20       polyvinylpyrrolidone, carboxymethylcellulose sodium, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, etc.

      The preferred isotonizing agent includes sodium chloride, glycerin, D-mannitol, etc.

- 25       The preferred buffering agent includes buffer solutions such as phosphate, acetate, carbonate, citrate, etc.

      The preferred local anesthetic includes benzyl alcohol, etc.

- 30       The preferred antiseptic includes p-hydroxybenzoic esters, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, sorbic acid, etc.

      The preferred antioxidant includes salts of sulfurous acid, ascorbic acid, etc.

- 35       The above pharmaceutical composition can be manufactured by conventional methods in the pharmaceutical

preparation techniques, for example methods described in the Japanese Pharmacopoeia.

5 Examples of dosage forms of the pharmaceutical composition include oral dosage forms such as tablets, capsules (inclusive of soft capsules and microcapsules), powders, granules, and syrups; and non-oral dosage forms such as injections, suppositories, pellets, and drip infusions. These dosage forms can be safely administered either orally or non-orally.

10 The dosage of the anti-inflammatory agent of the present invention differs depending on the subject, route of administration, clinical condition, etc. For oral administration to an adult patient, for instance, the usual unit dose is about 0.1 mg/kg to about 30 mg/kg, preferably  
15 about 2 mg/kg to about 20 mg/kg, as the compound of the invention which is an active ingredient, which dose is preferably administered once to 3 times a day.

#### BEST MODE FOR CARRYING OUT THE INVENTION

20 The following examples and test examples are intended to describe the present invention in further detail and should by no means be construed as defining the scope of the invention.

##### Example 1

25 A fluidized-bed granulating and drying machine (produced by powerex, Japan) was charged with 2479.5 g of hydrochloride of Compound No.1 (2250 g in terms of Compound No.1), 13930.5 g of lactose and 540 g of carboxymethylcellulose calcium (carmellose calcium),  
30 followed by mixing at the preheating temperature and spraying 7500 g of an aqueous solution containing 450 g of hydroxypropylcellulose to yield granules. 16820 g of the granules were processed with cutter-mill (produced by Showa Kagaku Kikai Kousakusho, Japan) to yield milled granules.  
35 16530 g of the milled granules, 513 g of carmellose calcium and 57 g of magnesium stearate were mixed to yield



lubricated powders by using tumbling mixer (produced by Showa Kagaku Kikai Kousakusho, Japan). 16800 g of the lubricated powders were tabletted by using tableting machine (produced by Kikusui Seisakusho, Japan) to yield  
 5 140000 tablets having the following formula and each containing 15 mg of Compound No. 1.

Formula per tablet (Unit: mg):

	1) Hydrochloride of Compound No.1	16.53
	2) Lactose	92.87
10	3) Carmellose calcium	7.2
	4) Hydroxypropylcellulose	3.0
	5) <u>Magnesium stearate</u>	<u>0.4</u>
	Total: 120.0	

#### 15 Example 2

In substantially the same manner as in Example 1, 140000 tablets having the following formula and each containing 30 mg of Compound No.1 were obtained.

Formula per tablet (Unit: mg):

20	1) Hydrochloride of Compound No.1	33.06
	2) Lactose	76.34
	3) Carmellose calcium	7.2
	4) Hydroxypropylcellulose	3.0
	5) <u>Magnesium stearate</u>	<u>0.4</u>
25	Total: 120.0	

#### Example 3

In substantially the same manner as in Example 2, 140000 tablets having the following formula and each  
 30 containing 45 mg of Compound No.1 were obtained.

Formula per tablet (Unit: mg):

	1) Hydrochloride of Compound No.1	49.59
	2) Lactose	114.51
	3) Carmellose calcium	10.8
35	4) Hydroxypropylcellulose	4.5
	5) <u>Magnesium stearate</u>	<u>0.6</u>

Total: 180.0

Test Example 1 (Reduction of plasma TNF- $\alpha$  level in mice)

The plasma TNF- $\alpha$  level was determined by using KKA<sup>y</sup> mice which are genetically obese, diabetic models, and a TNF- $\alpha$  inhibitory action of the compound of the present invention was evaluated.

Namely, eighteen male KKA<sup>y</sup> mice (10 week old), genetically obese, diabetic models, were divided into two groups each of which consists of nine mice. A powdered commercial diet (CE-2, produced by Japan Clea) was given to one group (control group), and the above powdered diet also containing 0.001 % (w/w) of hydrochloride of Compound No. 1 was given to the other group (drug administration group) ad libitum. Mice in these groups were bred for 4 days. The average dosage of drug per mouse was 16 mg/kg body weight/day. On the fourth day, mice were sacrificed and blood was collected in tubes containing heparin.

The collected blood was centrifuged and the plasma TNF- $\alpha$  level was determined by the enzyme immunoassay based on the biotin-streptavidin method. Namely, 5  $\mu$ l of a solution of an anti-TNF- $\alpha$  antibody IgG [produced by Genzyme, USA] (100  $\mu$ g/ml) diluted with 0.05 M Tris-HCl buffer (pH 8.0) was added to each wells of a 96-well polystyrene microtiter plate [produced by Falcon, USA], followed by standing at the room temperature for 2 hours to adhere the anti-TNF- $\alpha$  antibody IgG to the plate. After removal of an excess antibody solution, each wells was washed with 0.1 M Tris-HCl buffer (pH 7.6) containing 0.4 M NaCl, 0.1 % (w/w) bovine serum albumin, 0.1 % (w/w) NaN<sub>3</sub> and 1 mM MgCl<sub>2</sub> (hereafter referred to as a washing buffer).

Ten  $\mu$ l of plasma or standard solution of TNF- $\alpha$  [Serotec, Great Britain] was added to each wells, followed by standing for 2.5 hours at the room temperature. After each wells was washed with a washing buffer, 200  $\mu$ l of a solution of a biotinylated anti-TNF- $\alpha$  antibody IgG (35

ng/ml) diluted with a washing buffer was added, followed by standing over night at 4 °C. After each wells was washed with a washing buffer, 20 µl of a solution of a β-D-galactosidase-linked streptavidin [produced by Boehringer Mannheim GmbH, Germany] diluted 6000 fold with a washing buffer was added, followed by standing for one hour at the room temperature.

Then, each wells was washed with a washing buffer, and β-D-galactosidase activity of an immune complex fixed at a solid phase was assayed. Namely, 30 µl of a substrate [60 mM of 4-methylumbelliferyl-β-D-galactoside, produced by Sigma, USA] was added to each wells to start an enzyme reaction. After the reaction was conducted at the room temperature for 4 hours, the enzyme reaction was stopped by addition of 0.13 ml of 0.1 M glycine-NaOH buffer (pH 10.3).

The fluorescence intensity of the produced 4-methylumbelliferone was determined using a fluorescence spectrometer [Cyto Fluor II, PerSeptive Biosystems, USA] at the wavelengths of 350 and 460 nm for excitation and emission, respectively.

Then, the amount of TNF-α was calculated from the obtained fluorescence intensity using a separately prepared dose-response curve.

The results are shown in Table 1.

Table 1. Plasma TNF-α level (pg/ml)

Control group	Drug administration group (Present invention)
4.97±1.75	1.52±1.08**

Mean ± Standard Deviation; Significantly different from Control group (\*\*:p<0.01)

It is apparent from Table 1 that the compound of the present invention significantly reduced plasma TNF-α level in mice.

35

Test Example 2 (Reduction of plasma TNF-α level in rats)

The plasma TNF- $\alpha$  level was determined by using Wistar fatty rats which are genetically obese, diabetic models, and a TNF- $\alpha$  inhibitory action of the compound of the present invention was evaluated.

5 Namely, hydrochloride of Compound No. 1 was orally administered to sixteen male Wistar fatty rats (16 week old), genetically obese, diabetic models, via gastric tube at a dose of 3 mg/kg body weight/day. Ten rats were sacrificed before drug administration, and the first, second, third  
10 and fourth day after drug administration, respectively. Then, blood was collected.

As the normal group, ten Wistar lean rats (16 week old) were sacrificed without drug administration and blood was collected.

15 The collected blood was centrifuged, and the plasma TNF- $\alpha$  level was determined in substantially the same manner as in Test Example 1.

The results are shown in Table 2.

Table 2. Plasma TNF- $\alpha$  level (pg/ml)

	Days after drug administration	TNF- $\alpha$ level (pg/ml)
20	Normal group	0 56.9 $\pm$ 47.5
25	Control group	0 139.5 $\pm$ 50.0
	Present invention	1 109.9 $\pm$ 61.0 2 115.1 $\pm$ 59.0 3 69.9 $\pm$ 64.3 30 4 67.2 $\pm$ 70.6*

Mean  $\pm$  Standard Deviation; Significantly different from Control group (\*:p<0.05)

It is apparent from Table 2 that the compound of the present invention reduced the plasma TNF- $\alpha$  level in rats  
35 time-dependently.

Test Example 3 (Reduction of TNF- $\alpha$  content in skeletal muscle of rats)

The TNF- $\alpha$  content in skeletal muscle was determined by using Wistar fatty rats which are genetically obese, diabetic models, and a TNF- $\alpha$  inhibitory action of the compound of the present invention was evaluated.

Namely, hydrochloride of Compound No. 1 was administered to male Wistar fatty rats (16 week old), genetically obese, diabetic models in substantially the same manner as in Test Example 2. Ten rats were sacrificed before drug administration, and the first, second, third and fourth day after drug administration, respectively. Then, skeletal muscle was collected.

As the normal group, ten Wistar lean rats (16 week old) were sacrificed without drug administration and skeletal muscle was collected.

To the collected skeletal muscle, 0.1 M Tris-HCl buffer (pH 7.6) containing 1 M NaCl, 2 % (w/w) bovine serum albumin, 2 mM ethylenediaminetetraacetic acid disodium salt (EDTA), aprotinin (80 tripsin-inhibitory units/liter) and 0.02 % (w/w) NaN<sub>3</sub> was added in an amount of 20 weight times of the weight of the wet skeletal muscle. After ultrasonic disintegration, the mixture was centrifuged at 15000 rpm for 30 minutes to obtain a supernatant.

The amount of TNF- $\alpha$  in the obtained supernatant was determined in substantially the same manner as in Test Example 1.

The results are shown in Table 3.

Table 3. TNF- $\alpha$  content in skeletal muscle (pg/g wet weight)

	Days after drug administration	Amount of TNF- $\alpha$ (pg/g wet weight)
Normal group	0	156.7 $\pm$ 61.9
Control	0	356.6 $\pm$ 105.6

group		
Present	1	200.1±165.1*
invention	2	181.4±108.2**
	3	105.1± 96.4**
5	4	107.3± 95.7**

Mean ± Standard Deviation; Significantly different from Control group (\*:p<0.05, \*\*:p<0.01)

It is apparent from Table 3 that the compound of the present invention reduced the TNF- $\alpha$  content in skeletal muscle of rats significantly and almost time-dependently.

#### Test Example 4 (Suppression of the active oxygen production in neutrophils)

The in vitro effect of the compound of the present invention on suppression of the active oxygen production in neutrophils was evaluated by determining the amount of peroxides in cells.

Namely, venous blood was collected from male Wistar rats (6 week old) while adding heparin. To the collected blood, the same volume of an aqueous solution of 3 %(w/w) dextran was added for separation of blood cells. After the mixture was allowed to stand for 30 minutes, precipitates obtained by centrifugation was suspended with saline. The suspension was piled on Ficoll-Hypaque solution (Sigma, USA), followed by centrifugation.

From the obtained precipitates, erythrocytes were removed by hemolysis to separate neutrophils.

The hemolysis was conducted in the following manner. Namely, 4 ml of an ice-cooled 0.2 %(w/w) aqueous solution of NaCl was added to the above precipitates, which was suspended quickly, followed by standing for 20 to 30 seconds to puncture the erythrocytes. Then, 4 ml of an ice-cooled 1.6 %(w/w) aqueous solution of NaCl was added to the obtained suspension, which was mixed to yield a mixed solution having the same osmotic pressure with the erythrocytes before puncture. The mixed solution was

centrifuged at 4 °C at 150 × g for 5 minutes. After the supernatants were removed, the precipitates were washed with PBS (phosphate buffer saline).

The thus obtained erythrocytes were washed with saline, followed by addition of a minimum essential medium to prepare a neutrophils floating solution. The obtained neutrophils floating solution was fractionated into tubes so that the number of neutrophils per tube is 106.

Then, hydrochloride of Compound No. 1 or Compound No. 8 was added to the obtained tubes at the concentration of 1 μM. After incubation for one hour, a fluorescent pigment [DCFH-DA (2,7-dichlorofluoresceine diacetic acid)] was added, which was subjected to determination of the fluorescence intensity by FACScan (Becton Dickinson, USA).

As the control group, the fluorescence intensity in the case of adding no drug was determined.

The relative values of the fluorescence intensity in the drug addition group when the fluorescence intensity in the control group was 100 were calculated. These values were defined as the amount of peroxides caused by active oxygen derived from neutrophils.

The results are shown in Table 4.

Table 4. Fluorescence intensity and peroxide level

	Fluorescence intensity	Peroxide level
Control group	707	100
Hydrochloride of Compound No. 1 (Present invention)	466	66
Control group	377	100
Hydrochloride of Compound No. 8 (Present invention)	242	64

It is apparent from Table 4 that the compound of the present invention suppressed the active oxygen production

in neutrophils.

TNF- $\alpha$  is produced by various cells such as monocytes, macrophages, neutrophils, fibroblasts, epithelial cells, astrocytes, and etc. TNF- $\alpha$  increases production of active oxygen in neutrophils, which are suggested to have a close relation with occurrence of rheumatoid arthritis [Clinical and Experimental Rheumatology, vol. 15, pp.233-237 (1997); Inflammation, vol. 20, pp.427-438 (1996)].

Therefore, it is considered that the compound of the present invention exhibited suppressive effects on the active oxygen production by reducing TNF- $\alpha$  production or TNF- $\alpha$  sensitivity in neutrophils based on the results of Test Example 4.


#### Industrial Applicability

The anti-inflammatory agent of the present invention is used as an agent for prophylaxis and treatment of TNF- $\alpha$  mediated inflammatory diseases such as diabetic complications (e.g., retinopathy, nephropathy, neuropathy, disorders in the great arteries, etc.), rheumatoid arthritis, osteoarthritis of the spine, osteoarthritis, low back pain, gout, postoperative or traumatic inflammation, remission of swelling, neuralgia, sore throat, cystitis, hepatitis, pneumonia, and etc.



## CLAIMS

1. An anti-inflammatory agent which affects by way of a TNF- $\alpha$  inhibitory action and comprises a compound of the formula:



10


wherein R represents a hydrocarbon group that may be substituted or a heterocyclic group that may be substituted; Y represents a group of the formula -CO-, -CH(OH)-, or -NR<sup>3</sup>- where R<sup>3</sup> represents an alkyl group that  
 15 may be substituted; m is 0 or 1; n is 0, 1 or 2; X represents CH or N; A represents a chemical bond or a bivalent aliphatic hydrocarbon group having 1 to 7 carbon atoms; Q represents oxygen or sulfur; R<sup>1</sup> represents hydrogen or an alkyl group; ring E may have further 1 to 4 substituents, which may form  
 20 a ring in combination with R<sup>1</sup>; L and M respectively represent hydrogen or may be combined with each other to form a chemical bond; or a salt thereof.

2. An anti-inflammatory agent according to Claim 1, wherein the heterocyclic group represented by R is a 5- to  
 25 7-membered monocyclic and heterocyclic group containing 1 to 4 hetero-atoms selected from oxygen, sulfur and nitrogen in addition to carbon as ring members or a condensed heterocyclic group.

3. An anti-inflammatory agent according to Claim 1, wherein R represents a heterocyclic group that may be  
 30 substituted.

4. An anti-inflammatory agent according to Claim 3, wherein the heterocyclic group is pyridyl, oxazolyl or thiazolyl.


35 5. An anti-inflammatory agent according to Claim 1, wherein the partial structural formula:



- 5    6.    An anti-inflammatory agent according to Claim 1,  
wherein X represents CH.
7.    An anti-inflammatory agent according to Claim 1,  
wherein R<sup>1</sup> represents hydrogen.
8.    An anti-inflammatory agent according to Claim 1,
- 10    wherein L and M respectively represent hydrogen.
9.    An anti-inflammatory agent according to Claim 1,  
wherein the compound is 5-[4-[2-(5-ethyl-2-  
pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione.
10.    An anti-inflammatory agent according to Claim 1,
- 15    wherein the compound is (R)-(+)-5-[3-[4-[2-(2-furyl)-5-  
methyl-4-oxazolylmethoxy]-3-methoxyphenyl]propyl]-2,4-  
oxazolidinedione.
11.    Method for treating or preventing a TNF- $\alpha$  mediated  
inflammatory disease in a mammal in need thereof, which
- 20    comprises administering to said mammal an effective amount  
of a compound as defined in claim 1 or a pharmacologically  
acceptable salt thereof.
12.    Use of a compound as defined in claim 1 or a  
pharmacologically acceptable salt thereof for the
- 25    manufacture of an agent for prophylaxis or treatment of a  
TNF- $\alpha$  mediated inflammatory disease.

## ABSTRACT

An anti-inflammatory agent which affects by way of a  
 TNF- $\alpha$  inhibitory action and comprises a compound of the  
 formula:



wherein R represents a hydrocarbon group that may be  
 substituted or a heterocyclic group that may be  
 substituted; Y represents a group of the formula -CO-,  
 -CH(OH)-, or -NR<sup>3</sup>- where R<sup>3</sup> represents an alkyl group that  
 may be substituted; m is 0 or 1; n is 0, 1 or 2; X represents  
 CH or N; A represents a chemical bond or a bivalent aliphatic  
 hydrocarbon group having 1 to 7 carbon atoms; Q represents  
 oxygen or sulfur; R<sup>1</sup> represents hydrogen or an alkyl group;  
 ring E may have further 1 to 4 substituents, which may form  
 a ring in combination with R<sup>1</sup>; L and M respectively represent  
 hydrogen or may be combined with each other to form a  
 chemical bond; or a salt thereof.

(modified)

PTO/SB/106 (8-96)  
Approved for use through 9/30/98. OMB 0651-0032  
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

## Declaration and Power of Attorney For Patent Application

### 特許出願宣言書及び委任状

### Japanese Language Declaration

### 日本語宣言書

下記の氏名の発明者として、私は以下の通り宣言します。

As a below named inventor, I hereby declare that:

私の住所、私書箱、国籍は下記の私の氏名の後に記載された通りです。

My residence, post office address and citizenship are as stated next to my name.

下記の名称の発明に関して請求範囲に記載され、特許出願している発明内容について、私が最初かつ唯一の発明者（下記の氏名が一つの場合）もしくは最初かつ共同発明者であると（下記の名称が複数の場合）信じています。

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Anti-inflammatory Agent

上記発明の明細書（下記の欄でx印がついていない場合は、本書に添付）は、

the specification of which is attached hereto unless the following box is checked:

☐ 月 日に提出され、米国出願番号または特許協定条約国際出願番号を \_\_\_\_\_ とし、  
(該当する場合) \_\_\_\_\_ に訂正されました。

☒ was filed on August 20, 1998  
as United States Application Number or  
PCT International Application Number  
PCT/JP98/03692 and was amended on \_\_\_\_\_  
(if applicable).

私は、特許請求範囲を含む上記訂正後の明細書を検討し、内容を理解していることをここに表明します。

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

私は、連邦規則法典第37編第1条56項に定義されたとおり、特許資格の有無について重要な情報を開示する義務があることを認めます。

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

## Japanese Language Declaration

## (日本語宣言書)

私は、米国法典第35編119条(a)-(d)項又は365条(b)項に基づき下記の、米国外の国の少なくとも一カ国を指定している特許協力条約365(a)項に基づく国際出願、又は外国での特許出願もしくは発明者証の出願についての外国優先権をここに主張するとともに、優先権を主張している、本出願の前に出願された特許または発明者証の外国出願を以下に、枠内をマークすることで、示しています。

## Prior Foreign Application(s)

外国での先行出願

(Number)

(番号)

9-225302

(Country)

(国名)

Japan

(Day/Month/Year Filed)

(出願年月日)

21/08/1997

Priority Not Claimed

優先権主張なし

私は、第35編米国法典119条(e)項に基づいて下記の米国外の特許出願規定に記載された権利をここに主張いたします。

(Application No.)

(出願番号)

(Filing Date)

(出願日)

私は、下記の米国法典第35編120条に基づいて下記の米国外の特許出願に記載された権利、又は米国外を指定している特許協力条約365条(c)に基づき権利をここに主張します。また、本出願の各請求範囲の内容が米国法典第35編112条第1項又は特許協力条約で規定された方法で先行する米国外の特許出願に開示されていない限り、その先行米国外出願書提出日より本出願書の日本国内または特許協力条約国際提出日よりの期間中に入手された、連邦規則法典第37編1条56項で定義された特許資格の有無に関する重要な情報について開示義務があることを認識しています。

(Application No.)

(出願番号)

(Filing Date)

(出願日)

(Application No.)

(出願番号)

(Filing Date)

(出願日)

私は、私自身の知識に基づいて本宣言書中で私が行なう表明が真実であり、かつ私の入手した情報と私の信じていることに基づき表明が全て真実であると信じていること、さらに故意になされた虚偽の表明及びそれと同等の行為は米国法典第18編第1001条に基づき、罰金または拘禁、もしくはその両方により処罰されること、そしてそのような故意による虚偽の声明を行なえば、出願した、又は既に許可された特許の有効性が失われることを認識し、よってここに上記のごとく宣誓を致します。

I hereby claim foreign priority under Title 35, United States Code, Section 119 (a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

(Application No.)

(出願番号)

(Filing Date)

(出願日)

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s), or 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of application.

(Status: Patented, Pending, Abandoned)

(現況: 特許許可済、係属中、放棄済)

(Status: Patented, Pending, Abandoned)

(現況: 特許許可済、係属中、放棄済)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

(modified)

PTO/SB/106 (8-96)  
Approved for use through 9/30/98. OMB 0651-0032  
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

### Japanese Language Declaration (日本語宣言書)

委任状: 私は下記の発明者として、本出願に関する一切の手続きを米特許商標局に対して遂行する弁理士または代理人として、下記の者を指名いたします。(弁護士、または代理人の氏名及び登録番号を明記のこと)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith (list name and registration number)

Philippe Y. RIESEN (Reg. No. 35,657) Miriam SOHN (Reg. No. 35,368)

Send Correspondence to:

書類送付先

Philippe Y. Riesen  
Intellectual Property Department  
Takeda Chemical Industries, Ltd.  
17-85, Jusohonmachi 2-chome  
Yodogawa-ku, Osaka 532-8686 Japan

Direct Telephone Calls to: (name and telephone number)

直接電話連絡先: (名前及び電話番号)

Philippe Y. Riesen  
(Phone) 81-6-6300-6124 (Japan)  
(Fax) 81-6-6300-6601  
(E-mail) riesen\_philippe@takeda.co.jp

唯一または第一発明者名		Full name of sole or first inventor	
ODAKA Hiroyuki			
発明者の署名	日付	Inventor's signature	Date
		<i>Hiroyuki Odaoka</i>	Nov 5, 1999
住所		Residence	
12-12, Katsuragi 2-chome, Kita-ku, Kobe-shi, HYOGO 651-1223 Japan JPX			
国籍		Citizenship	
Japan			
私書箱 Takeda Chemical Industries (IPD)		Post Office Address	
17-85, Jusohonmachi 2-chome, Yodogawa-ku, Osaka 532-8686 Japan			
第二共同発明者		Full name of second joint inventor, if any	
MOMOSE Yu			
第二共同発明者	日付	Second inventor's signature	Date
		<i>Yu Momose</i>	Nov 5, 1999
住所		Residence	
8-7, Sumiregaoka 2-chome, Takarazuka-shi, HYOGO 665-0847 Japan JPX			
国籍		Citizenship	
Japan			
私書箱 Takeda Chemical Industries (IPD)		Post Office Address	
17-85, Jusohonmachi 2-chome, Yodogawa-ku, Osaka 532-8686 Japan			

(第三以降の共同発明者についても同様に記載し、署名をすること)

(Supply similar information and signature for third and subsequent joint inventors.)